



Seven decades of Ramadan intermittent fasting research: Bibliometrics analysis, global trends, and future directions



Khaled Obaideen ^a, Katia H. Abu Shihab ^b, Mohamed I. Madkour ^c, MoezAllIslam E. Faris ^{b, *}

^a Sustainable Energy & Power Systems Research Centre, RISE, University of Sharjah, P.O. Box 27272, Sharjah, United Arab Emirates

^b Department of Clinical Nutrition and Dietetics, University of Sharjah, Sharjah, United Arab Emirates

^c Department of Medical Laboratory Sciences, University of Sharjah, Sharjah, United Arab Emirates

ARTICLE INFO

Article history:

Received 6 June 2022

Received in revised form

28 June 2022

Accepted 3 July 2022

Keywords:

Artificial intelligence

Bibliometric analysis

Citation analysis

Medical journals

Scientometric

Text mining

ABSTRACT

Background and aims: There is a large body of research focused on various aspects related to Ramadan intermittent fasting (RIF) and human health and disease. This study aimed to quantify the bibliometric data of RIF medical research over the past seven decades and explore these variables qualitatively via text mining analysis.

Methods: We used the Scopus search engine to identify published articles related to RIF from inception to December 31, 2021. All types of research articles were included. Scientometric and bibliometric measures were determined using Excel, Biblioshiny, and VOSviewer. This study proposed a bibliometric and text mining method to qualitatively and quantitatively recognize the RIF research trend.

Results: The Scopus search returned 1915 relevant articles. Most citations pertained to publications from the last two decades, and most publications were original research articles. These publications had received around 27,000 citations, and the 20 most prolific publishing journals had an average h-index of 112.25. More than one-third of all medical publications were in open-access journals. There was a 13-fold increase in medical research on RIF over the past few decades. We identified the 10 most prolific publishing countries, institutes, journals, and authors. We also identified five scientific hotspots of RIF scientific literature, which were: diabetes, metabolic health, public health, physiology, and maternity.

Conclusion: This is the first comprehensive bibliometric analysis of medical research related to RIF. The research gaps identified will shape future research directions and foster collaborative research activities toward enhanced medical nutrition research revolving around RIF.

© 2022 Diabetes India. Published by Elsevier Ltd. All rights reserved.

1. Background

Currently, Muslims constitute the world's second-largest religious group. There are an estimated 2.0 billion Muslims, accounting for about 25% of the 8 billion global population [1]. Every year, the Islamic holy month of Ramadan is observed by more than 1.5 billion Muslims worldwide [2]. During Ramadan, Muslims must refrain from eating, drinking, smoking, and sexual activity from dawn to dusk each day, for about 11–22 hours depending on the geographical location and solar season. This holy month is also a dedicated time for Muslims to practice humility and increase their acts of worship. Ramadan has attracted extensive research attention over the years because of its impact on human health, especially given its unique intermittent fasting (IF) model.

Intermittent fasting (IF) is linked with numerous positive health effects such as the prevention of metabolic, inflammatory, and degenerative diseases [3–7]. Different regimens are present for IF, including time-restricted feeding, alternate-day fasting, and religious forms, such as Ramadan IF (RIF) [4]. A growing body of evidence, including original research and meta-analyses, supports the disease-preventing and health-improving effects of RIF on healthy and overweight/obese adults. These effects include reducing body weight [8], changing body composition [9], with emphasis on lowering visceral adiposity [10], improving liver functions [11], ameliorating the metabolic syndrome components, improving cardiometabolic risk factors [12], normalizing glucometabolic markers [8], and alleviating inflammatory and oxidative stress markers [13–15]. The drastic shifts in dietary and lifestyle habits accompanying the changes in circadian rhythm [16] sleep quality [17], and food selections [18], are also concomitant with changes in the expressions of multiple genes regulating circadian rhythms such as the *CLOCK* gene [19], antioxidant and metabolism-

Abbreviations: RIF, Ramadan intermittent fasting.

* Corresponding author.

E-mail addresses: mfaris@sharjah.ac.ae, moezfaris@hotmail.com (M.E. Faris).

controlling genes (*SIRT1*, *SIRT3*, *TFAM*, *SOD2*, and *Nrf2*) [20] and fat mass and obesity-associated protein gene *FTO* [21].

Assessing existing literature on RIF may offer clear directions for further research on Ramadan. To date, there has been no comprehensive quantitative and qualitative analysis of existing research on RIF in the general context of health. Therefore, this study explored existing scientific literature on RIF and its impact on health via a bibliometric analysis and text mining. Bibliometrics is a systemic statistical analysis of existing literature in a given field, its general scientific nature, and its development over time [22]. Such analysis sheds light on the most prolific research countries, institutions, and authors, as well as the most cited relevant publications. Bibliometrics can be coupled with text mining, which refers to the analysis and modeling of unstructured natural language text present in literature through machine learning and linguistics [23]. Text mining can help in processing a large amount of textual data from a scientific database, creating text clusters, and extracting meaningful patterns or concepts for the overall existing body of scientific literature. The key objectives of this study were to analyze quantitative and qualitative trends in Ramadan medical research, including: annual growth rates; numbers of publications and citations; top contributing authors, institutions, and countries; highest publishing journals for research on RIF; and thematic research hotspots on RIF in terms of health. This comprehensive analysis will provide insights for future researchers on medical and health topics related to RIF that have been explored and highlight any gaps that remain to be addressed. Such an enhanced scientific approach to RIF may therefore offer a potential turning point for the health of those who observe this holy month.

2. Materials and methods

2.1. Design

The present study used general bibliometric methods for a quantitative analysis of the scientific output of the literature on RIF. This was complemented by text mining, which provided a qualitative outlook on the research patterns of RIF, thematic fields, and top publishing authors and countries. The methodology and analysis were based on several previous bibliometric studies [23–27].

2.2. Data collection and search strategy

The bibliographic database on RIF research was extracted from the Scopus platform [28], on December 31, 2021. Scopus provides a comprehensive database of source-neutral abstracts and citations, with over 25,100 titles from more than 5000 international publishers in different research fields [29]. Fig. 1 presents the data extraction and analysis steps. The search strategy used for this study (Fig. 2) included the word “Ramadan” and different spellings, such as “Ramadhan” and “Ramazan.” We also included other phrases that are commonly used to refer to the Ramadan month of fasting, such as “Islamic fasting,” “diurnal fasting,” “consecutive 30 days of fasting,” and “religious fasting.” The search query template (TITLE-ABS-KEY) offered by Scopus was used to search these terms in the titles, abstracts, and keywords of publications in the database. To restrict the search to publications related to RIF and health, we excluded non-medical subject areas, such as social sciences, arts and humanities, agricultural sciences, environmental sciences,




Fig. 1. Workflow steps of search query formulation for data extraction (flow figure A) and different methods of data analysis (flow figure B).

```

(TITLE-ABS-KEY (ramadan) OR TITLE-ABS-KEY (ramadhan) OR TITLE-ABS-KEY (ramazan) OR TITLE-ABS-KEY (
"Islamic fasting") OR TITLE-ABS-KEY ("diurnal fasting") OR TITLE-ABS-KEY ("Ramadan intermittent fasting") OR TITLE-ABS-
KEY ("Ramadan diurnal intermittent fasting") OR TITLE-ABS-KEY ("consecutive 30 days of fasting") OR TITLE-ABS-KEY (
"religious fasting")) AND (EXCLUDE (SUBJAREA, "ARTS") OR EXCLUDE (SUBJAREA, "BUSI") OR EXCLUDE (
SUBJAREA, "CENG") OR EXCLUDE (SUBJAREA, "CHEM") OR EXCLUDE (SUBJAREA, "COMP") OR EXCLUDE (
SUBJAREA, "EART") OR EXCLUDE (SUBJAREA, "ECON") OR EXCLUDE (SUBJAREA, "ENGI") OR EXCLUDE (
SUBJAREA, "ENVI") OR EXCLUDE (SUBJAREA, "MATE") OR EXCLUDE (SUBJAREA, "MATH") OR EXCLUDE (
SUBJAREA, "PHYS") OR EXCLUDE (SUBJAREA, "SOCI"))

```

Fig. 2. Search query used for searching for medical (health) research on Ramadan fasting in the Scopus database, that includes 'Ramadan fasting' synonyms and excludes non-medical fields.

mathematics, engineering, economics, computer sciences, physics, chemistry, earth, and planetary sciences, decision sciences, and other similar fields. We did not restrict the time frame, and the scope of the publication years ranged from 1952 to 2021, with the first publication on RIF in the Scopus database dating from 1952. The search returned 1915 publications.

2.3. Data management and statistical analysis

The Scopus database of relevant publications was extracted and analyzed using the Bibliometrix package (Biblioshiny software) in R [30]; Biblioshiny is an open-source comprehensive tool that integrates several different science mapping analysis techniques. The extracted data was converted into a bibliometrix file through Biblioshiny for further analysis and chart creation in Microsoft Excel 2016. The quantitative parameters analyzed included yearly output, document type, number of citations, publication language, the most cited publications, and the most prolific countries, institutions, journals, and authors. The measurements were ranked using Standard Competition Ranking (SCR), as used in other bibliometric publications. The quality of the presented journals was assessed by the SCLMago Journal Rank (SJR). Although Scopus provides sophisticated algorithmic profiling of its publishing authors, not all publications are accurately matched to a single author profile if an author has different abbreviations for their name. This results in authors seemingly having fewer publications on the Scopus platform than they actually do, as their publications are dispersed among their different author profiles. For example, abbreviations for Alawadi F. (10 Ramadan-related publications): Alawadi F. (one publication), Al awadi F. (two publications), Al awadi FF. (two publications) all pertain to a single author (Al-Awadi, Fathehya Fardallah; 15 publications) on Scopus. Therefore, a thesaurus of all author name abbreviations and the number of relevant publications was extracted using VOSviewer software [31] and used to merge author names by cross-checking their respective Scopus profiles. VOSviewer is software that allows the creation, visualization, and text-mining of different bibliometric variables. This helped in obtaining a more accurate report of the number of publications contributed by each author in the Scopus database. To identify the themes covered by the most prolific authors, the top keywords in their publications were identified using SciVal which is functionality for in-depth analysis offered by Scopus.

In addition, text mining was performed using the VOSviewer software, and is presented in collaboration or co-occurrence networks for different parameters (authors, countries, and index words) with different analytical visual overlays. To create a map of diverse thematic hotspots of RIF research, the top 500 index words

(those with at least 16 occurrences) were used as a unit of analysis; however, 461 index words were included in the final map after selective filtering. Generic and non-thematic index words (e.g., human, male, female, age, article, and others), country names, and journal names were excluded to provide a clear presentation of research hotspots. Different spellings of medical terminology (e.g., hypoglycemia and hypoglycemia) were merged using the VOSviewer thesaurus functionality. In the resulting map, the distance between the index words indicated the strength of co-occurrence in the same publications. Collaboration network maps of the most prolific authors (those with at least 15 publications) and countries (those with at least 25 publications) were also created [32]. Color-coded clusters of all maps were generated using VOSviewer, with a resolution level of 1.00. Finally, the world map charts were created using Datawrapper [33], a web-based mapping tool, based on the data from the Scopus analysis results.

3. Results

3.1. Document type, yearly output, publication language, and citations

Globally, there were 1915 medical research publications on RIF over the last seven decades, with the first publication from 1952 and the majority (97%, $n = 1865$) of publications during the last three decades (1989–2021) (Fig. 3). The peak of citations was for publications from 1975, but the majority of citations (85%, 22,565 citations) pertained to publications from the last two decades (1997–2019). Fig. 4 presents an abstract assortment of the general bibliometric nature of the publications, such as the total number of extracted publications, their total citations, approximate publishing authors, top 3 prolific publishing journals, and percentage of articles with open access. It additionally presents the distribution of the publications' dominant languages, document type, Scopus research fields, and a brief overview of the publications' thematic hotspots. On date of the publication search, 35% of the publications on Ramadan were with open-access (OA) journals.

3.2. Authors and sources

Ramadan research increased over the last three decades from about 10 publications per year during the 1990s to an average of 130 publications for 2020/2021. As shown in Fig. 5, RIF has been extensively researched all over the world with publications from 90 countries/regions as of as of publications extraction date. The most contributed publications were from the UK and Saudi Arabia (around 250 and 200 publications, respectively). This was followed




Fig. 3. Annual publication and citation frequency of medical Ramadan research. The majority (97%, $n=1865$) of publications are of the last three decades (1989–2021), while the bulk of citations (85%, $n=22,565$) pertained to publications from the last two decades (1997–2019).




Fig. 4. Abstract assortment of general bibliometric qualities of medical publications on Ramadan fasting. The publications experienced 8.9% annual growth rate, and pertained to around 10 Scopus fields, primarily Medicine. 35% of publications were open-access.

by the US and Iran (around 180 publications each), Türkiye (around 160 publications), Tunisia (around 150 publications), and the

United Arab Emirates (around 130 publications).

International co-authorship networks were identified in four




Fig. 5. Two comparative color-coded world maps of countries that published Ramadan research (Map B) and the percentage of the Muslim population in those countries in 2021 (Map A) [107]. Despite the majority of Muslims globally residing in Middle East/North Africa region (Map B), the UK was leading in Ramadan publications, followed by Saudi Arabia, US, and Iran (Map A).

clusters, as presented in Fig. 6. The analysis is based upon the frequency of co-authorship between Ramadan researchers and represented in terms of the authors' respective affiliated countries. In map A of Fig. 6, Several clusters appeared to consist of regionally close countries, particularly the green- and blue-colored clusters.

The green cluster comprised countries located in South/South-east Asia, along with the geographically nearby Australia. This cluster also contained Türkiye, which is located in West Asia/Middle East and is a part of the Mediterranean region: it was strongly linked to the other Middle Eastern countries in the blue cluster. Similarly, the




Fig. 6. Co-authorship network of 30 affiliated countries over the last seven decades (1952–2021). In map A, the size of the circles and their links reflect the publication frequency and the strength of their connections respectively. The countries of the same color belong to the same collaboration cluster, which reflects the frequent collaboration of the authors of the same publications affiliated with these countries. The countries are grouped into 4 clusters, depicting the influence of geographical proximity on authors collaboration. In map B, the presented countries are color-coded on a spectrum from violet = near 2012 to yellow = near 2016, according to their average publication year. In map C, the presented countries are color-coded on a spectrum from white = near 10 citations to violet = near 25 citations, according to the average citations received per publication. The size of the presented country's respective circles reflects their total citations received. In map D, the presented countries are color-coded on a spectrum reflecting their average normalized received citations per publication from pale yellow = near 1.0 normalized citations to red-orange = near 1.6 normalized citations. The size of the circles reflect the country's total received normalized citations.

blue cluster mostly included the countries bordering Saudi Arabia. The red cluster contained regionally close countries surrounding the Mediterranean Sea, including Italy, Spain, Tunisia, and France. Iran-affiliated authors appeared to have the highest co-authorship solely with US-affiliated authors, which was represented by the strong link between the two countries. US-, UK-, Saudi Arabia-, and UAE-affiliated authors appeared to have the highest international co-authorships, as they collaborated with 28, 29, 28, and 27 countries respectively out of the 30 countries presented in the figure. US-affiliated authors published a minimum of 10 articles in collaboration with at least one country from each presented cluster.

Map B of Fig. 6 adds a visual overlay of the average publication years for the aforementioned countries, which mostly ranged from 2012 to 2016, as analyzed by VOSviewer software. The majority of publications from Canada, the United Arab Emirates (UAE), Bahrain, Italy, Indonesia, and Bangladesh were relatively recent compared with those from the UK, Iran, and Saudi Arabia, which were the highest publishing countries. Map C of Fig. 6 shows an overlay of average citations received per publication. France, Morocco, Kuwait, Lebanon, and Algeria had the highest average citations per publication, with Kuwait having the most with an average of 31 citations per publication on average. Most other countries had 15–20 citations per publication on average. To correct for the fact that older documents had had more time to receive citations, the data were normalized in VOSviewer and presented in map D of Fig. 6. After

normalization, the UAE, Bahrain, Jordan, Kuwait, and Italy had the highest citations per publication annually, with the UAE and Bahrain leading at around 1.9 citations per publication per year.

Tehran University of Medical Sciences (Iran), King Saud University (Saudi Arabia), and Dubai health authority (UAE) were the top three institutions of Ramadan-related research (Table 1). Table 2 details the top 15 journals that published on RIF from 1952 to 2021, along with their citations, citation/article ratio, SJR (SCImago Journal Rank), and h-Index. *Diabetes Research and Clinical Practice*, *Saudi Medical Journal*, *Nutrients*, *Journal of Sport Sciences*, *Tunisiaie Medicale*, and *PLoS ONE* were the top five journals for publishing Ramadan research. However, on a citation basis, *Diabetes Research and Clinical Practice* (1074 citations), followed by *Journal of Sports Sciences* (591 citations).

Among the top 20 cited publications, diabetes management during Ramadan was the most cited topic, followed by the impact of Ramadan on human health and its effects on body weight and metabolism. Other highly cited research fields included the impact of Ramadan on human circadian rhythms, cardiometabolic profile, and athletic physical performance. The most cited publications were the EPIDIAR study [34], a diabetes management paper [35], and a study on metabolic and weight changes during Ramadan [36] (Table 3).

The top 17 authors (top 10 by SCR) with the most publications

Table 1

Top 10 prolific institutions in terms of published Ramadan-related medical research over the last seven decades (1952–2021).

SCR ^a	Affiliation	Articles
1	Tehran University of Medical Sciences (Iran)	68
2	King Saud University (Saudi Arabia)	66
3	Dubai Health Authority (UAE)	47
4	Baqai Medical University (Pakistan)	37
5	Kuwait University (Kuwait)	32
6	University of Genoa (Italy)	31
7	University of Sharjah (UAE)	31
8	Shahid Beheshti University of Medical Sciences (Iran)	30
9	Cairo University (Egypt)	29
10	Universiti Putra Malaysia (Malaysia)	27

^a SCR, Standard Competition Ranking (ranking in which equal institutions were given the same ranking number).

focused on Ramadan are presented in Table 4. The most prolific authors were Hassanein M., Chtourou H., and Chamari K. with 51, 42, and 34 Ramadan-related publications, respectively. An analysis of the top keywords used in the publications showed that the majority of the top prolific authors explored Ramadan in the context of diabetes or athletic performance. Fewer authors were interested in the relationships between RIF and circadian rhythms and weight change. The collaboration networks of the top 25 authors with at least 13 publications are presented in Fig. 7's map A, with five color-coded clusters indicative of frequent collaboration among specific authors. From the authors' respective affiliations, it could be inferred that most authors' clusters were based locally. For example, the yellow author cluster (Chtourou H., Aloui A., and Souissi N.) in Tunisia and the red cluster (Hassanein M., Hanif W., Ghouri N., Chowdhury TA., and Beshyah SA.) were predominantly affiliated with the UK, the UAE, or both. The greatest extent of international collaboration was seen among authors in the green cluster, as they were affiliated with an array of countries (e.g., Chamari K.: Qatar, Tunisia, Ireland; Chaouachi A.: Vietnam, New Zealand, Tunisia; Aziz AR.: Singapore). Due to the different abbreviations of the author names as mentioned in the methods and materials above, there might be uncertainty in the authors'

publications numbers.

Fig. 7's map B adds an overlay of the average production year. Most of the top prolific authors had their Ramadan-related publications clustering around 2010 to 2018. The majority of these authors had recent publications (Faris ME., Bragazzi NL., Ahmedani MY., Ghouri N., Al-awadi FF., and Kalra S.). In addition, six authors (Hassanein M., Chowdhury TA., Beshyah SA., Hanif W., Chtourou H., and Bahammam AS.) tended to publish around 2015–2016. This corresponded to the exponential growth in publications over the last three decades.

4. Research hotspots and categories

Fig. 8's map A showed that the index words for Ramadan research appeared to be clustered into five research fields that studied its impact on diabetes (blue cluster), public health (red cluster), metabolic profile (green cluster), human physiology (orange cluster), and maternity (purple cluster). Table 5 presents the top index words for each cluster.

4.1. Diabetes

Diabetes is one of the most thoroughly researched diseases in Ramadan literature, as its publications' index words were visualized as a whole individual cluster. Most publications appear to focus on diabetes mellitus type 2 (313 occurrences), complications of hypoglycemia (289 occurrences), and hemoglobin A1C or HbA1c (200 occurrences) as a glucose parameter. Diabetes management including the adjustment of hypoglycemic agents (175 occurrences), insulin (242 occurrences), metformin (110 occurrences), and gliclazide (75 occurrences) were common keywords in this cluster. Ramadan research also focused on the efficacy of patient education (83 occurrences) including blood glucose monitoring (109 occurrences) and patient compliance (67 occurrences).

4.2. Metabolic profile

This cluster reflects generally publications addressing metabolism (145 occurrences), as Ramadan fasting (373 occurrences)

Table 2

Top 20 publishing journals on Ramadan medical research over the last seven decades (1952–2021).

SCR ^a	Source	Articles		Citations	C/A ^b	SJR ^c	H-index
		N	%				
1	Diabetes Research and Clinical Practice	77	4.02	906	11.77	1.605	114
2	Saudi Medical Journal	33	1.72	511	15.48	0.437	51
3	Nutrients	25	1.30	284	11.36	1.418	115
4	Journal of Sports Sciences	24	1.25	591	24.63	1.208	137
5	PLoS ONE	23	1.20	142	6.17	0.99	332
5	Tunisie Medicale	23	1.20	400	17.39	0.199	18
6	Pakistan Journal of Medical Sciences	21	1.10	149	7.10	0.316	30
7	Eastern Mediterranean Health Journal	20	1.04	250	12.50	0.442	47
8	Annals of Saudi Medicine	19	0.99	391	20.58	0.373	44
8	The British Medical Journal	19	0.99	312	16.42	1.831	429
9	International Journal of Clinical Practice	18	0.94	105	5.83	0.756	98
9	Practical Diabetes International	18	0.94	420	23.33	0.172 ^a	0
10	Biological Rhythm Research	16	0.84	211	13.19	0.252	33
11	Diabetes and Metabolic Syndrome: Clinical Research and Reviews	15	0.78	580	38.67	0.684	40
12	Diabetic Medicine	15	0.78	95	6.33	1.474	145
13	Asian Journal of Sports Medicine	14	0.73	144	10.29	0.216	27
13	British Journal of Nutrition	14	0.73	311	22.21	1.073	118
13	Journal of the Pakistan Medical Association	14	0.73	589	42.07	0.242	43
15	Diabetes Care	13	0.68	1074	82.62	6.636	363
16	Singapore Medical Journal	12	0.63	545	45.42	0.452	61

^a SCR: Standard Competition Ranking (ranking in which equal journals were given the same ranking number).

^b C/A, citations and articles ratio (citation frequency divided by the article frequency for each journal).

^c SJR, SCImago Journal Rank.

Table 3

Top cited articles on Ramadan medical research over the last seven decades.

SCR ^a	Authors	Title	Citations	Journal	Document type
1	Salti et al. [34]	A population-based study of diabetes and its characteristics during the fasting month of Ramadan in 13 countries: Results of the Epidemiology of Diabetes and Ramadan 1422/2001 (EPIDIAR) study	450	Diabetes Care	Article
2	Al-Arouj et al. [35]	Recommendations for management of diabetes during Ramadan: Update 2010	241	Diabetes Care	Note
3	Ziaee et al. [36]	The changes in metabolic profile and weight during Ramadan fasting	192	Singapore Medical Journal	Article
4	Trepanowski et al. [95]	The impact of religious fasting on human health	190	Nutrition Journal	Review
5	El Ati et al. [96]	Increased fat oxidation during Ramadan fasting in healthy women: An adaptative mechanism for body-weight maintenance	189	American Journal of Clinical Nutrition	Article
6	Al-Arouj et al. [97]	Recommendations for management of diabetes during Ramadan	176	Diabetes Care	Review
6	Leiper et al. [98]	Effects on health of fluid restriction during fasting in Ramadan	176	European Journal of Clinical Nutrition	Article
7	Bogdan et al. [99]	Ramadan fasting alters endocrine and neuroendocrine circadian patterns. Meal-time as a synchronizer in humans?	168	Life Sciences	Article
8	Adlouni et al. [100]	Fasting during Ramadan induces a marked increase in high-density lipoprotein cholesterol and a decrease in low-density lipoprotein cholesterol	157	Annals of Nutrition and Metabolism	Article
9	Bravis et al. [101]	Ramadan education and awareness in diabetes (read) programme for Muslims with type 2 diabetes who fast during Ramadan	156	Diabetic Medicine	Article
9	Roky et al. [102]	Physiological and chronobiological changes during Ramadan intermittent fasting	156	Annals of Nutrition and Metabolism	Review
10	Hassanein et al. [103]	Diabetes and Ramadan: Practical guidelines	152	Diabetes Research and Clinical Practice	Review
11	Trepanowski et al. [57]	Impact of caloric and dietary restriction regimens on markers of health and longevity in humans and animals: A summary of available findings	139	Nutrition Journal	Review
12	Roky et al. [104]	Sleep during Ramadan intermittent fasting	129	Journal of Sleep Research	Article
13	Faris et al. [15]	Intermittent fasting during Ramadan attenuates proinflammatory cytokines and immune cells in healthy subjects	126	Nutrition Research	Article
15	Sweileh et al. [105]	Body composition and energy metabolism in resting and exercising Muslims during Ramadan fast	124	Journal of Sports Medicine and Physical Fitness	Article
16	Iraki et al. [106]	Ramadan diet restrictions modify the circadian time structure in humans. A study on plasma gastrin, insulin, glucose, and calcium and on gastric pH	120	Journal of Clinical Endocrinology and Metabolism	Article

^a SCR: Standard Competition Ranking (ranking in which equal journals were given the same ranking number).

was strongly linked to several keywords: blood glucose level (389 occurrences), blood pressure (167 occurrences), blood triacylglycerol level (144 occurrences), blood cholesterol level (141 occurrences), and body weight (353 occurrences). A good amount of studies focused on RIF's impact on pathophysiology (135 occurrences), and possibly on metabolic diseases, such as obesity (44 occurrences) and metabolic syndrome (18 occurrences). Furthermore, the impact of RIF on inflammatory markers [37,38] (e.g., interleukin-6: 20 occurrences, c-reactive protein: 28 occurrences) and oxidative stress (24 occurrences) were also evident in Ramadan research. Indicators of kidney function (43 occurrences), such as blood creatinine level (74 occurrences), blood albumin level (26 occurrences), blood potassium levels (34 occurrences), and glomerular filtration rate (24 occurrences) were also comprehensively explored, particularly in fasting patients with kidney disease (11 occurrences) or kidney failure (32 occurrences). Blood parameters that may affect the development of kidney stones and change during RIF, such as blood uric acid levels (77 occurrences) were also frequently researched topics.

4.3. Human physiology

Ramadan research was also directed toward the effects of diurnal intermittent dietary abstention and changes in food habits (120 occurrences) during Ramadan on human physiology (227 occurrences) in different biological aspects. An area of particular interest was the change in athletic performance (49 occurrences) in fasting Muslim athletes (51 occurrences) during Ramadan, as shown in this cluster. Most studies explored fasting athletes'

adaptation (31 occurrences) and (optimization of) exercise (103 occurrences) to maintain their usual physical performance (29 occurrences) during Ramadan [39]. Ramadan research also addressed the changes in body clock or circadian rhythm (114 occurrences) related to fasting, which is strictly time-specific (keyword: time, 113 occurrences) from pre-dawn to dusk. This key term was strongly linked to body weight due to its metabolic associations [40]. As circadian rhythms are also highly affected by sleep (73 occurrences), sleep time (41 occurrences), fatigue (41 occurrences), and somnolence (20 occurrences) were frequently addressed by Ramadan researchers. In addition, Ramadan research intensively explored changes in dietary consumption among fasting Muslims, particularly energy, carbohydrates, and fluid intakes (134, 70, and 46 occurrences, respectively), physical activity (116 occurrences), and overall body composition (69 occurrences) during Ramadan. Finally, Ramadan research also frequently targeted the topic of dehydration (106 occurrences) during RIF, which was shown to be strongly linked to diabetes-related keywords.

4.4. Public health

Ramadan research was also clustered around the relationship between Ramadan and public health (20 occurrences), as reflected by several keywords: disease severity (77 occurrences), disease duration (65 occurrences), disease incidence (74 occurrences), hospitalization (68 occurrences), and prevalence (55 occurrences). Many of the published papers focused on the epidemiology, particularly of diabetes mellitus (238 occurrences), which was shown by its strong link to the keyword "risk factors" (144

Table 4

Top authors with the highest number of publications on Ramadan medical research over the last seven decades.

SCR ^a	Authors	Articles		Citations ^d	Countries of affiliation	Top three key-phrases ^e
		(authors' name variations merged ^b)	In Scopus ^c			
1	Hassanein M.	51	45	1495	UK, UAE	Insulin and non-insulin-dependent diabetes; Diabetes
2	Chtourou H.	42	40	628	Tunisia	Soccer player; Athletes; Athletic performance
3	Chamari K.	34	33	861	Tunisia, France, Italy, Qatar, Ireland	Athletic performance; Anaerobic exercise; Soccer player
4	Bahammam AS.	29	10	594	Canada, KSA	Sleep; Disorders of excessive somnolence; Circadian rhythm
5	Bragazzi NL.	28	25	290	Italy, France, Russia, Canada, Morocco, Palestine,	Systemic review; Athletes; Chronic kidney failure
5	Souissi N.	28	27	600	France, Tunisia	Soccer player; Athletic performance; Circadian rhythm
6	Chaouachi A.	23	23	671	Tunisia, Vietnam, New Zealand	Soccer player; Athletic performance; Time of day
7	Trabelsi K.	21	19	222	Tunisia, Bahrain, Austria	Strength training; Sleep pattern; Dietary carbohydrates.
8	Beshyah S.	19	19	255	UK, UAE	Hypoglycemia; Diabetes; Non-insulin-dependent diabetes
8	Aloüi A.	19	18	229	Tunisia	Soccer player; Athletic performance; Physical capacity
8	Ahmedani MY.	19	16	189	Norway, Pakistan	Diabetes; Eating habits
9	Faris ME.	17	11	333	Jordan, KSA, UAE	Caloric restriction; Meta-analysis; Weight control
9	Aziz AR.	17	15	399	Singapore	Athletes; Athletic performance; Fitness
10	Kalra S.	16	16	68	India, Canada, Pakistan	India; Patient care; Diabetes
10	Hanif W.	15	16	505	UK	Diabetes; Non-insulin-dependent diabetes; Insulin replacement therapy
10	Bener A.	15	15	402	US, UK, KSA, Kuwait, UAE, India, Türkiye, Qatar	NA
10	Al-Awadi FF.	15	10	121	UAE	Diabetes type 2; Continuous glucose monitoring; Diabetes

^a SCR, Standard Competition Ranking (ranking in which authors with an equal number of publications were given the same ranking number).^b Several authors had different variations of how their names appeared in the Scopus database. These names with their respective publications were merged to calculate the accurate number of publications for each author.^c Number of publications presented by the available analysis functionality in Scopus.^d Number of citations corresponding to the merged number of publications for the respective author.^e Top key phrases identified via the Scopus SciVal function. This corresponds to the author's publications number identified by the Scopus artificial intelligence and was limited to 2011–2021. SciVal was not able to identify the keywords used by Bener A.

occurrences) in this cluster. The lifestyle changes (98 occurrences) that accompany RIF were also of interest in Ramadan epidemiological studies, which may explain the observed changes in health parameters. This was reflected by its strong links to the keywords “body weight,” “blood glucose level,” and “diabetes mellitus type 2” in the other two clusters. The effects of Ramadan on Muslims' psychological status (92 occurrences) and mental health were also a focal point of this cluster. Finally, Ramadan studies also explored Muslim patients' attitude (44 occurrences) toward health education (34 occurrences), and the cultural competence of healthcare personnel (65 occurrences) to properly care for fasting Muslim patients during Ramadan, especially in predominantly non-Muslim countries [41].

4.5. Maternity

Ramadan studies also considered the impact of RIF on pregnancy outcomes (130 occurrences), such as newborns' birth weight (25 occurrences). Pregnancy was also strongly linked to diabetes-related keywords such as blood glucose level, diabetes mellitus type 1, and diabetes mellitus type 2.

In terms of citations, there were publications with particular index words that received more citations on average than others in the same cluster, as presented in Fig. 8's map C after normalization for publication year. For example, in the diabetes cluster, the index words “diabetic medications” (e.g., metformin and sulfonylurea) and “side effects” (e.g., headache, vomiting, nausea, diarrhea, weight gain) attracted the most citations. Index words indicative of lipid profile (e.g., triglycerides, lipids), inflammatory status (e.g., oxidative stress, interleukin 6, inflammation), body weight change (e.g., weight loss, obesity, body mass index) were the most prominent hotspots in the metabolic profile cluster. In the human

physiology cluster, most index words received a relatively high number of citations annually on average, especially index words related to dietary habits (e.g., energy intake, carbohydrate intake, protein intake) and circadian rhythms. In the public health cluster, the index words “COVID-19” and “epidemiology” received the highest normalized average citations. An overlay of the publication year, presented in Fig. 8's map B, shows that most of these clusters were thoroughly explored from the start of the last decade onwards. Index words that addressed inflammatory status and susceptibility to develop cancer (e.g., tumor necrosis factor, oxidative stress, interleukin 6, inflammation) were some of the commonly used words in recent research on RIF. In addition, words supportive of the current COVID-19 pandemic (e.g., COVID-19, adverse events, healthcare personnel) were also relatively recent researched fields.

5. Discussion

According to our novel findings, over the past seven decades, research attention towards Ramadan has increased in the last three decades. This corresponds with the rising growth rate of Muslim populations around the world. From 1990 to 2010, an average annual growth rate of 2.2% in Muslim populations globally was reported [42]. Additionally, OA publications on Ramadan steeply increased over the last two decades from 16% in 2000–2006 to around 50% in 2016–2021. Governmental support facilitated OA publishing in many Arab and Islamic countries as part of the endeavor to improve the ranking of universities in those countries [43,44].

UK, Saudi Arabia, the US, and Iran were the highest RIF research contributors, and their affiliated authors had the highest international co-authorship. Nonetheless, our extracted publications are limited to the ones published in the English language, which can




Fig. 7. Collaboration networks of the 25 most prolific authors with at least 13 publications. In map A, the authors with the same color belong to the same collaboration cluster, which reflects their frequent collaboration on the same publications. There are five collaboration clusters; most are based locally according to the authors' respective country affiliations. In map B, the circles of the presented authors are color-coded on a spectrum from dark blue = near 2010 to yellow = near 2018, according to their average year of publication.

skew the representation of the international collaborations. The international collaborations seem to be affected by the geographical proximity of the researchers, as seen in a few of several co-authorship clusters in Fig. 6. Many France-affiliated authors had particularly strong links with authors from Tunisia, Morocco, and Canada, which may attribute to French being the second most common language of Ramadan-related publications. This points to the role of language in determining the extent of international collaboration [45]. Nonetheless, the variation in fasting daytime hours [46], Muslim ethnicities, and cultures across the world call for more collaborative controlled studies to allow better assessment of the impact of Ramadan.

Our analysis of top-cited articles, top prolific journals, and the top prolific authors' research interests on Ramadan shows that the scientific interest in RIF is predominantly around diabetes, metabolic health, and athletic performance. This coincides with the five identified thematic hotspots (diabetes, metabolic health, human physiology, public health, and maternity) and reflects the growing attention to diverse research on human health during Ramadan, especially as Muslims are one of the most prominent religious populations worldwide.

Diabetes is the top researched disease in the Ramadan literature, as the high occurrence of its relevant keywords constructed an individual thematic cluster on its own, unlike the other researched




Fig. 8. Thematic clusters of the top index words in Ramadan medical research over the last seven decades. In map A, the size of the circles and their links reflect the frequency and strength of the connections, respectively. Index words with the same color belong to the same thematic cluster, which reflects their co-occurrences in the same publications. Ramadan research clusters into five thematic hotspots: metabolic profile, physiology, diabetes, public health, and maternity. In map B, the circles of the words are color-coded on a spectrum from dark blue = near 2010 to yellow = near 2018, according to the average year of publication. In map C, the circles of the index words are color-coded on a spectrum from white = near 0.6 normalized citations to red-orange = near 2.0 normalized citations, according to the average normalized citations received.

diseases. Diabetes is also the research interest for almost a third of the identified top prolific authors on Ramadan. With urbanization, population growth, and population aging, diabetes mellitus has continued to rise as a global health burden [47]. RIF observance has no adverse effects on glucose homeostasis markers in healthy people, yet the situation differs for patients with diabetes [8]. RIF may impact glucose blood levels and increase the risk for hypoglycemia among people with diabetes [8]. Therefore, Islamic law allows unwell individuals for whom fasting may aggravate their condition to be exempt from fasting [48]. However, 79% of Muslims with type 2 diabetes still fasted during Ramadan, as noted in an epidemiological study involving 13 Muslim countries (EPIDIAR) in 2004 [34]. This motivated the International Diabetes Federation in collaboration with Ramadan and Diabetes International Alliance to formulate and release special practical guidelines on Ramadan and diabetes in 2021–2022 [49]. The possible improvements in metabolic status observed during RIF [50], which is also highly researched as elaborated below, may have attributed to the intensive research on diabetes.

RIF's potential to improve body composition and metabolism has been explored in many Ramadan studies [8,48,51,52], as seen in the metabolic profile cluster. There's a global steep rise in other metabolic diseases, like obesity [53], and likewise, metabolic syndrome [54] which is a collection of risk factors associated with diabetes or cardiovascular disease development [55,56]. Therefore, the changes in metabolic biomarkers [57], body weight [58], the risk of metabolic diseases [59], and overall metabolism experienced by fasting Muslims during Ramadan were also of high research interest. Some Ramadan publications also address RIF's effects on inflammatory markers and oxidative [37,38].

Additionally, Ramadan involves changes in sleeping habits [17], eating behaviors, food selections [18], and increased acts of spirituality [60,61]. This formed the basis for the human physiology thematic cluster of the analyzed Ramadan literature. Many identified prolific authors on RIF consistently published on the physiological changes experienced by fasting Muslim athletes during

Ramadan. It was surprising not to find the term “chrononutrition”. Chrononutrition is an emerging scientific field that refers to the interplay between nutrition, timed eating patterns, and metabolic health [62]. Given the strict timed nature of RIF, chrononutrition has potential for research in the scope of Ramadan and may be interrelated with both metabolic health, wellness, and sleep quality [16]. Several authors had a particular interest in Ramadan's impact on circadian rhythm changes, which are affected by the changes in eating and sleeping timings of fasting Muslims [63].

These aforementioned physiological and health changes may have attributed to the scientific interest in disease severity, disease prevalence, and hospitalization [64] during Ramadan which is evident in the public health thematic cluster. Additionally, many Muslims during Ramadan adopt an active nocturnal lifestyle, causing a significant shift in emergency department visits from daytime to nighttime [65]. As Ramadan involves avoiding immorality, practicing self-control, and focusing on increased acts of worship as well as dietary abstention, relatively fewer publications also explored Ramadan on Muslims' mental health [66].

Although pregnant and lactating women are religiously exempt from fasting, many women opt to fast during Ramadan for personal or cultural reasons [67,68]. Therefore, many studies exist on RIF's impact on pregnancy and neonatal outcomes as seen in the maternity thematic cluster. The links existing between the words 'pregnancy' and diabetes-related terms could be due to common complications of glucose intolerance experienced during pregnancy, i.e. gestational diabetes mellitus [69].

Overall, the fields of diabetes and metabolic disease in the context of RIF have been thoroughly addressed from many different perspectives compared with the other hotspots. Ramadan's impact on human physiology and public health also appeared to have been intensively studied, although not as expansively. Another neglected area of research is the study of mental health, especially in the context of spirituality during Ramadan [70].

Insufficient literature exists on RIF's impact on patients with chronic illnesses who opt to observe Ramadan. These include

Table 5

Top 10 indexed words for each thematic cluster of Ramadan medical research over the last seven decades.

Cluster	Top 10 occurring index words	Occurrences	Average citations	Average normalized citations	Average publication year
METABOLIC PROFILE (GREEN CLUSTER)	Glucose blood level	389	18.5	1.4	2012.9
	Body weight	353	22.2	1.6	2013.3
	Blood pressure	167	17.6	1.2	2013.6
	Metabolism	145	19.8	1.7	2013.3
	Triacylglycerol level	142	26.8	1.6	2011.5
	Cholesterol blood level	141	26.4	1.4	2010.9
	HDL cholesterol	141	27.3	1.6	2012.3
	Pathophysiology	135	14.8	1.4	2015.9
	LDL cholesterol	131	27.0	1.7	2012.9
	Lipids	84	26.5	1.8	2011.5
	Overall	4345	20.6	1.5	2013.4
PHYSIOLOGY (YELLOW CLUSTER)	Physiology	227	15.4	1.4	2014.6
	Energy intake	134	30.4	1.8	2011.7
	Food habits	120	27.5	1.5	2010.1
	Physical activity	116	22.1	1.6	2014.7
	Circadian rhythm	114	30.5	1.7	2010.0
	Time	113	28.3	1.7	2011.0
	Dehydration	106	21.9	1.3	2011.7
	Exercise	103	27.2	1.6	2012.4
	Feeding behavior	80	28.6	1.6	2009.1
	Sleep	73	30.9	1.7	2011.5
	Overall	4309	26.4	1.5	2011.7
DIABETES (BLUE CLUSTER)	Diabetes mellitus type 2	313	17.9	1.4	2014.1
	Hypoglycemia	289	18.2	1.4	2014.5
	Insulin aspart	242	20.1	1.4	2013.4
	Hemoglobin a1c	200	16.6	1.4	2014.9
	Hypoglycemic agents	175	20.3	1.5	2014.3
	Diabetes mellitus type 1	139	18.1	1.3	2014.3
	Glycemic control	115	11.1	1.2	2016.7
	Hyperglycemia	115	18.6	1.4	2014.7
	Metformin	110	22.3	1.7	2014.6
	Sulfonylurea	109	24.0	1.8	2014.2
	Overall	4224	22.4	1.7	2013.8
PUBLIC HEALTH ^a (RED CLUSTER)	Diabetes mellitus	238	15.3	1.1	2012.2
	Risk factors	144	16.2	1.4	2013.8
	Lifestyle	98	21.7	1.6	2013.0
	Risk assessment	93	20.3	1.4	2013.7
	Psychology	92	9.7	0.8	2014.4
	Religion and medicine	83	16.4	1.0	2008.1
	Child	78	11.2	0.9	2011.5
	Disease severity	77	10.5	0.9	2014.4
	Incidence	74	21.3	1.4	2014.1
	Adverse event	68	5.7	0.9	2018.1
	Overall	3743	13.3	1.1	2013.4
MATERNITY (PURPLE CLUSTER)	Pregnancy	130	16.6	1.0	2011.1
	Body height	32	10.8	0.9	2015.7
	Gestational age	32	16.3	0.9	2011.8
	Birth weight	25	20.4	1.2	2010.0
	Newborn	25	18.1	1.2	2011.3
	Maternal nutrition	23	12.7	0.8	2011.7
	Pregnancy diabetes mellitus	20	13.1	1.5	2017.6
	Echography	17	6.8	0.6	2013.6
	Pregnancy outcome	15	14.6	0.9	2010.4
	Lactation	14	31.0	1.0	2004.5
	Overall	496	16.0	1.0	2012.2

^a Generic words (Ramadan fasting, adult, Islam, Muslim) were omitted from the top occurring public health index words list and overall cluster parameter calculations as they were not specifically relevant.

cancer, autoimmune and inflammatory bowel diseases, mental health problems, and patients with immune-compromised diseases such as HIV/AIDS who self-decide to observe RIF. Future RIF research must be directed toward clinical and basic studies in different health and disease conditions other than nutrition and metabolic disorders [71].

6. Future directions

With the innovation of omics techniques that can elaborate on the biological changes accompanying different human life

processes, it is rational to apply this multi-disciplinary approach in studying the various changes accompanying RIF. Among these, genomics, proteomics, metabolomics, metagenomics, and transcriptomics are the main omics of interest. Applying these techniques in studying RIF will help in collectively characterizing and quantifying the pools of biological molecules that are released during fasting. This will expand our understanding of the RIF's effect on different health and disease conditions and will guide in the decision-making process when caring for their Muslim fasting patients during Ramadan. In a non-RIF model, intermittent prolonged fasting showed significant changes in more than 370

metabolites, including plasma amino acids, glycerol, free fatty acids, and ketone bodies which reflect changes in the body's fuel management under fasting conditions [72]. However, only one study showed the various metabolomic changes that accompanied the practice of RIF [73]. Our lab work on the metabolomic and lipidomic changes in RIF revealed variable changes in the serum metabolites pertaining to caffeine, amino acids, fatty acid metabolism, and different ceramides and sphingolipids after starting RIF compared with pre-fasting levels (unpublished data) [108].

At the proteomic level, examining the impact of intermittent fasting [74], LC–MS/MS analysis of human plasma for participants on an 8-week intermittent fasting regimen revealed that this pattern was associated with significant changes in different apolipoproteins, involved in lipoprotein metabolism, with a significant decrease in plasma triglycerides after intermittent fasting. In the context of RIF, however, Mindikoglu and colleagues published two important works that showed RIF triggered an anticancer serum proteome response [75] and upregulated key regulatory proteins for circadian rhythms, lipid and glucose metabolism, circadian remodeling, cytoskeleton remodeling, DNA repair, the immune system and cognitive function in healthy subjects. Further research is warranted to explore the other metabolic and inflammatory markers related to RIF, and apply other omics techniques (genomics, epigenomics, lipidomics, and transcriptomics) that may be affected by the dietary and lifestyle modifications associated with RIF.

The human gut microbiome and its fundamental role in human health and disease, as well as its ability to modulate variable biochemical and physiological changes, have been well established [76,77]. Therefore, it is important to examine the changes in gut microbiota during RIF and elaborate on their metabolic and inflammatory impacts, especially considering the various dietary and lifestyle modifications accompanying RIF. Recently, few studies examining the effect of RIF on human gut microbiota have been published, which showed significant improvements and changes in different types of microbial flora [78–81].

Artificial intelligence (AI), and its multiple forms and applications (including machine learning and neural networks), are the most distinguishing features of the current and coming eras. They can offer essential tools that could be applied in examining the relationship between the practice of RIF and outcomes related to health and disease. One paper applied AI to patients with type 2 diabetes for the first time in an attempt to predict the occurrence of hypoglycemic events during their fasting [82]. More extensive application of AI, however, is warranted to help other patient populations, such as tools to assist patients with kidney and gastrointestinal diseases who opt to observe RIF.

With the globalization of the COVID-19 pandemic and the role of RIF in the yearly worship of Muslims, it is necessary to elaborate on the impact of observing RIF on innate and adaptive immunity among various population groups, especially sensitive and high-risk groups such as pregnant and lactating women, children, and older people who opt to fast during Ramadan, especially given the increased susceptibility to COVID-19 infection when fasting. Of the few studies that investigated the impact of observing intermittent fasting on immunity, autophagy, and other related mechanisms, two reports revealed that intermittent fasting could be used as a priming tool for host defense against SARS-CoV-2 infection by modulating the autophagy and immune response [83,84]. An important review presented the interplay of intermittent fasting, obesity, and COVID-19 infection, and showed how fasting mitigated the obesity-induced predisposing factors that lowered the efficacy of immunity against infection [85]. For RIF and COVID-19, few studies discussed the plausible impacts of observing RIF on immunity. The first study was a narrative review by Faris and

colleagues [86], which reported the expected beneficial effect of observing RIF based on the published literature concerning the immunity-related factors of RIF. This work was followed by a few articles showing the same expected beneficial effect of RIF on immunity against SARS-CoV-2 infection [87–89].

7. Strengths and limitations

This is the first comprehensive bibliometric study that quantitatively and qualitatively analyzed the medical RIF literature. The analysis was based upon Scopus database, a multidisciplinary database that covers a broad spectrum of universal journals. Scopus covers 57% of citations from six databases (Scopus, Microsoft Academic, Google Scholar, Web of Science (WoS), Dimensions, and OpenCitations) [90]. In addition, Scopus covers a larger number of documents that are not covered by CWTS WoS, including documents with substantial numbers of citations and references [91].

There was no time restriction upon publications extracted, therefore our findings are representative of the quantitative, thematic, and temporal trends of Ramadan research over the last seven decades. Finally, the extracted database underwent exhaustive data cleaning to enhance the accuracy of the data presentation, particularly in terms of authors' publications and thematic hotspots of Ramadan scientific literature. This study was limited by lack of inclusion of scientific journals that were not indexed in Scopus. Although most authors tend to publish in international journals, some choose to publish in emerging journals out of institutional or national loyalty [92,93]. This might have affected our representation's accuracy of the available literature on Ramadan fasting, and the extent of authors' works and collaborations. Our study was able to partially compare the results with similar bibliometric work on Ramadan fasting in the scope of diabetes, although many differences were found because of the generality of our scope and restriction only to the Scopus database [27]. Therefore, future bibliometric work on Ramadan literature should consider including more than one database.

8. Conclusion

This study presents a comprehensive quantitative and qualitative overview of the scientific productivity of RIF research over the last seven decades. It identified the RIF research publication trends and its most prolific countries, journals, institutions, and authors with their scientific interests. The complementary text mining identified these authors' and countries' collaboration networks and productivity trends. It also identified the most researched themes of RIF, along with their temporal trends and citation frequency. Our study demonstrated the growing scientific attention toward RIF. Nonetheless, there's an oversaturation of RIF research on diabetes, metabolic health, and physiology. Therefore, our findings will assist researchers in determining the gaps in knowledge about RIF and can also contribute to future collaborations and research directions.

Ethics approval

Not Applicable.

Consent for publication

Not Applicable.

Availability of data and materials

The datasets analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors report there are no competing interests to declare.

Funding

This research received no specific grant from any funding agency, commercial or not-for-profit sectors.

Author contribution: CRediT author statement

Conceptualization, KO, MF; Methodology, KO; Software, KO; Validation, KO, KA; Formal analysis, KO, KA; Investigation, MF, KO, KA; Resources, MF, KO; Data Curation, KO, KA; Writing - Original Draft, KA, KO, MF; Writing - Review & Editing, MF, MM; Visualization, KA, KO; Supervision, MF; Project administration, MF, KO.

Acknowledgments

Deep thanks are due to the librarian Mr. Nadeem Rafiq and the library staff, Central Library of the University of Sharjah, for the continuous support in providing the full-text articles requested.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.dsx.2022.102566>.

References

- [1] WorldPopulationReview. Muslim population by country 2022. 2022.
- [2] Abdelrahim D, Faris ME, Hassanein M, Shakir AZ, Yusuf AM, Almeneessier AS, et al. Impact of ramadan diurnal intermittent fasting on hypoglycemic events in patients with type 2 diabetes: a systematic review of randomized controlled trials and observational studies. *Front Endocrinol* 2021;12.
- [3] Mattson MP, Longo VD, Harvie M. Impact of intermittent fasting on health and disease processes. *Ageing Res Rev* 2017;39:46–58.
- [4] Patterson RE, Sears DD. Metabolic effects of intermittent fasting. *Annu Rev Nutr* 2017;37:373–93.
- [5] De Cabo R, Mattson MP. Effects of intermittent fasting on health, aging, and disease. *N Engl J Med* 2019;381:2541–51.
- [6] Di Francesco A, Di Germanio C, Bernier M, De Cabo R. A time to fast. *Science* 2018;362:770–5.
- [7] Zeb F, Wu X, Chen L, Fatima S, Haq I-u, Chen A, Majeed F, Feng Q, Li M. Effect of time-restricted feeding on metabolic risk and circadian rhythm associated with gut microbiome in healthy males. *Br J Nutr* 2020;123:1216–26.
- [8] Faris ME, Jahrami H, BaHammam A, Kalaji Z, Madkour M, Hassanein M. A systematic review, meta-analysis, and meta-regression of the impact of diurnal intermittent fasting during Ramadan on glucometabolic markers in healthy subjects. *Diabetes Res Clin Pract* 2020;165:108226.
- [9] Fernando H, Zibellini J, Harris R, Seimon R, Sainsbury A. Effect of ramadan fasting on weight and body composition in healthy non-athlete adults: a systematic review and meta-analysis. *Nutrients* 2019;11:478.
- [10] Faris ME, Madkour MI, Obaideen AK, Dalah EZ, Hasan HA, Radwan H, et al. Effect of Ramadan diurnal fasting on visceral adiposity and serum adipokines in overweight and obese individuals. *Diabetes Res Clin Pract* 2019;153:166–75.
- [11] Faris ME, Jahrami HA, Abdelrahim DN, Bragazzi N, BaHammam AS. The effects of Ramadan intermittent fasting on liver function in healthy adults: a systematic review, meta-analysis, and meta-regression. *Diabetes Res Clin Pract* 2021;178.
- [12] Jahrami HA, Faris ME, Janahi AI, Janahi MI, Abdelrahim DN, Madkour MI, et al. Does four-week consecutive, dawn-to-sunset intermittent fasting during Ramadan affect cardiometabolic risk factors in healthy adults? A systematic review, meta-analysis, and meta-regression. *Nutr Metabol Cardiovasc Dis* 2021;31:2273–301.
- [13] Faris ME, Jahrami HA, Obaideen AA, Madkour MI. Impact of diurnal intermittent fasting during Ramadan on inflammatory and oxidative stress markers in healthy people: systematic review and meta-analysis. *J Nut Intermediary Metabol* 2019;15:18–26.
- [14] Faris ME, Hussein RN, Al-Kurd RA, Al-Farajeh MA, Bustanji YK, Mohammad MK. Impact of Ramadan intermittent fasting on oxidative stress measured by urinary 15-isoprostane. *J Nutri Metabol* 2012;2012.
- [15] Faris ME, Kacimi S, Al-Kurd RA, Farajeh MA, Bustanji YK, Mohammad MK, et al. Intermittent fasting during Ramadan attenuates proinflammatory cytokines and immune cells in healthy subjects. *Nutr Res* 2012;32:947–55.
- [16] BaHammam AS, Almeneessier AS. Recent evidence on the impact of Ramadan diurnal intermittent fasting, mealtime, and circadian rhythm on cardiometabolic risk: a review. *Front Nutr* 2020;7:28.
- [17] Faris ME, Jahrami HA, Alhayki FA, Alkhawaja NA, Ali AM, Aljeeb SH, et al. Effect of diurnal fasting on sleep during Ramadan: a systematic review and meta-analysis. *Sleep Breath* 2020;24:771–82.
- [18] Shatila H, Baroudi M, El Sayed Ahmad R, Chehab R, Forman MR, Abbas N, Faris M, Naja F. Impact of ramadan fasting on dietary intakes among healthy adults: a year-round comparative study. *Front Nutr* 2021;8. 689788–689788.
- [19] Ajabnoor GMA, Bahijri S, Shaik NA, Borai A, Alamoudi AA, Al-Aama JY, Chrousos GP. Ramadan fasting in Saudi Arabia is associated with altered expression of CLOCK, DUSP and IL-1alpha genes, as well as changes in cardiometabolic risk factors. *PLoS One* 2017;12:e0174342.
- [20] Madkour MI ATE-S, Jahrami HA, Sherif NM, Hassan RE, Awadallah S, Faris ME. Ramadan diurnal intermittent fasting modulates SOD2, TFAM, Nrf2, and sirtuins (SIRT1, SIRT3) gene expressions in subjects with overweight and obesity. *Diabetes Res Clin Pract* 2019;155:107801.
- [21] Madkour M, Abdel-Rahman W, Malhab L, Abdel-Rahman W, Abdelrahim D, Saber-Ayad M, et al. Ramadan diurnal intermittent fasting is associated with attenuated FTO gene expression in subjects with obesity: a prospective cohort study. *Front Nutr* 2022.
- [22] Wallin JA. Bibliometric methods: pitfalls and possibilities. *Basic Clin Pharmacol Toxicol* 2005;97:261–75.
- [23] Talabis M, McPherson R, Miyamoto I, Martin J, Kaye D. Analytics defined. *Information Security Analytics*; 2015. p. 1–12.
- [24] Ho L, Alonso A, Forio MAE, Vanclooster M, Goethals PL. Water research in support of the Sustainable Development Goal 6: a case study in Belgium. *J Clean Prod* 2020;277:124082.
- [25] Kalantari A, Kamsin A, Kamaruddin HS, Ebrahim NA, Gani A, Ebrahimi A, et al. A bibliometric approach to tracking big data research trends. *J. Big Data* 2017;4:1–18.
- [26] Song G, Wu J, Wang S. Text mining in management research: a bibliometric analysis. *Secur Commun Network* 2021;2021.
- [27] Beshyah WS, Beshyah SA. Bibliometric analysis of the literature on Ramadan fasting and diabetes in the past three decades (1989–2018). *Diabetes Res Clin Pract* 2019;151:313–22.
- [28] MacKellar M, Vigerust DJ. Role of haptoglobin in health and disease: a focus on diabetes. *Clin Diabetes*: a publication of the American Diabetes Association 2016;34:148–57.
- [29] Elsevier S. Scopus content coverage guide. Amsterdam: Elsevier BV; 2020.
- [30] Aria M, Cuccurullo C. bibliometrix: an R-tool for comprehensive science mapping analysis. *J informet* 2017;11:959–75.
- [31] Eck NJv, Waltman L, Vos. A new method for visualizing similarities between objects. In: *Advances in data analysis*. Springer; 2007. p. 299–306.
- [32] Waltman NJvEaL. VOSviewer Manual. version 1.6.17 edition22. July 2021.
- [33] Datawrapper. Create charts and maps [software]. <https://www.datawrapper.de/>.
- [34] Salti I, Bénard E, Detournay B, Bianchi-Biscay M, Le Brigand C, Voinet C, Jabbar A, Group ES. A population-based study of diabetes and its characteristics during the fasting month of Ramadan in 13 countries: results of the epidemiology of diabetes and Ramadan 1422/2001 (EPIDIAR) study. *Diabetes Care* 2004;27:2306–11.
- [35] Al-Arouj M, Assaad-Khalil S, Buse J, Fahdil I, Fahmy M, Hafez S, Hassanein M, Ibrahim MA, Kendall D, Kishawi S. Recommendations for management of diabetes during Ramadan: update 2010. *Diabetes Care* 2010;33:1895–902.
- [36] Ziaee V, Razaee M, Ahmadinejad Z, Shaikh H, Yousefi R, Yarmohammadi L, Bozorgi F, Behjati MJ. The changes of metabolic profile and weight during Ramadan fasting. *Singap Med J* 2006;47:409.
- [37] Faris ME, Kacimi S, Refat A, Farajeh MA, Bustanji YK, Mohammad MK, et al. Intermittent fasting during Ramadan attenuates proinflammatory cytokines and immune cells in healthy subjects. *Nutr Res (NY)* 2012;32:947–55.
- [38] Mo'ez Al-Islam EF, Jahrami HA, Obaideen AA, Madkour MI. Impact of diurnal intermittent fasting during Ramadan on inflammatory and oxidative stress markers in healthy people: systematic review and meta-analysis. *J Nut Intermediary Metabol* 2019;15:18–26.
- [39] Maughan RJ, Zerguini Y, Chalabi H, Dvorak J. Achieving optimum sports performance during Ramadan: some practical recommendations. *J Sports Sci* 2012;30:S109–17.
- [40] Kiriyaama K, Yamamoto M, Kim D, Sun S, Yamamoto H, Oda H. Skipping breakfast regimen induces an increase in body weight and a decrease in muscle weight with a shifted circadian rhythm in peripheral tissues of mice. *Br J Nutr* 1:26.
- [41] Abolaban H, Al Moujahed A. Muslim patients in Ramadan: a review for primary care physicians. *Avicenna journal of medicine* 2017;7:81–7.
- [42] Lugo L, Cooperman A, O'Connell E, Stencil S. The future of the global Muslim population. USA: Pew Research Center; 2011.
- [43] Khallaf AM. Library and information science open access journals in Arab countries. 2017.
- [44] Boufarrs M. Charting the Open Access scholarly journals landscape in the UAE. *Scientometrics* 2020;122:1707–25.
- [45] Yao B. In: *International research collaboration: challenges and opportunities*, vol. 37. Los Angeles, CA: SAGE Publications Sage CA; 2021. p. 107–8. 107–108.
- [46] Azahhaf N, Ramadan. Longest and shortest fasting hours globally. In:

- Morocco world news: q; 2021; 2021.
- [47] Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care* 2004;27: 1047–53.
 - [48] Jaleel MA, Raza SA, Fathima FN, Jaleel BNF. Ramadan and diabetes: as-saum (the fasting). *Ind J Endocrinol Metabol* 2011;15:268.
 - [49] Hassanein M, Afandi B, Ahmedani MY, Alamoudi RM, Alawadi F, Bajaj HS, Basit A, Bennakhi A, El Sayed AA, Hamdy O. Diabetes and ramadan: practical guidelines 2021. *Diabetes Res Clin Pract* 2022;109185.
 - [50] Tahapary DL, Astrella C, Kristanti M, Harbuwono DS, Soewondo P. The impact of Ramadan fasting on metabolic profile among type 2 diabetes mellitus patients: a meta-analysis. *Diabetes Metabol Syndr: Clin Res Rev* 2020;14: 1559–70.
 - [51] Faris ME, Jahrami H, Abdelrahman D, Bragazzi N, BaHammam A. The effects of Ramadan intermittent fasting on liver function in healthy adults: a systematic review, meta-analysis, and meta-regression. *Diabetes Res Clin Pract* 2021;178:108951.
 - [52] Jahrami HA, Alsibai J, Obaideen AA, Faris ME. Impact of Ramadan diurnal intermittent fasting on the metabolic syndrome components in healthy, non-athletic Muslim people aged over 15 years: a systematic review and meta-analysis. *Br J Nutr* 2020;123:1–22.
 - [53] Jaacks LM, Vandevijvere S, Pan A, McGowan CJ, Wallace C, Imamura F, Mozaffarian D, Swinburn B, Ezzati M. The obesity transition: stages of the global epidemic. *Lancet Diabetes Endocrinol* 2019;7:231–40.
 - [54] Saklayen MG. The global epidemic of the metabolic syndrome. *Curr Hypertens Rep* 2018;20:1–8.
 - [55] Oguoma VM, Coffee NT, Alsharrah S, Abu-Farha M, Al-Refaei FH, Alkandari A, Al-Mulla F, Daniel M. Anthropometric cut-points for discriminating diabetes and the metabolic syndrome among Arabs and Asians: the Kuwait Diabetes Epidemiology Program. *Br J Nutr* 2022;127:92–102.
 - [56] Mirmiran P, Aghayan M, Bakhshi B, Hosseinpour-Niazi S, Azizi F. Socioeconomic status and lifestyle factors modifies the association between snack foods intake and incidence of metabolic syndrome. *Nutr J* 2021;20:1–12.
 - [57] Trepanowski JF, Canale RE, Marshall KE, Kabir MM, Bloomer RJ. Impact of caloric and dietary restriction regimens on markers of health and longevity in humans and animals: a summary of available findings. *Nutr J* 2011;10:107.
 - [58] Bakhotmah BA. The puzzle of self-reported weight gain in a month of fasting (Ramadan) among a cohort of Saudi families in Jeddah, Western Saudi Arabia. *Nutr J* 2011;10:1–8.
 - [59] Nematy M, Alinezhad-Namaghi M, Rashed MM, Mozhdehifard M, Sajjadi SS, Akhlaghi S, Sabery M, Mohajeri SAR, Shalaei N, Moohebaty M. Effects of Ramadan fasting on cardiovascular risk factors: a prospective observational study. *Nutr J* 2012;11:1–7.
 - [60] Pathy R, Mills KE, Gazeley S, Ridgley A, Kiran T. Health is a spiritual thing: perspectives of health care professionals and female Somali and Bangladeshi women on the health impacts of fasting during Ramadan. *Ethn Health* 2011;16:43–56.
 - [61] Ali Khan MM, Nor NM, Mamat NM, Mohd-Shukri NA. Abu bakar wam: fasting in Islam: a combination of spiritual elevation and prevention of diseases. *IJUM Med J Malaysia* 2018;17.
 - [62] Flanagan A, Bechtold DA, Pot GK, Johnston JD. Chrono-nutrition: from molecular and neuronal mechanisms to human epidemiology and timed feeding patterns. *J Neurochem* 2021;157:53–72.
 - [63] Qasrawi SO, Pandi-Perumal SR, BaHammam AS. The effect of intermittent fasting during Ramadan on sleep, sleepiness, cognitive function, and circadian rhythm. *Sleep Breath* 2017;21:577–86.
 - [64] AlZahrani AM, Zawawi MM, Almutairi NA, Alansari AY, Bargawi AA. The impact of Ramadan on visits related to diabetes emergencies at a tertiary care center. *BMC Emerg Med* 2021;21:1–10.
 - [65] Butt T, Khan HU, Ahmed I, Eldali A. Emergency department attendance patterns during Ramadan. *Ann Saudi Med* 2016;36:258–64.
 - [66] Koushali AN, Hajiamini Z, Ebadi A, Bayat N, Khamseh F. Effect of Ramadan fasting on emotional reactions in nurses. *Iran J Nurs Midwifery Res* 2013;18: 232.
 - [67] Faris ME, Al-Holy MA. Implications of Ramadan intermittent fasting on maternal and fetal health and nutritional status: a review. *Mediterr J Nutr Metabol* 2014;7:107–18.
 - [68] Kridli SA-O. Health beliefs and practices of Muslim women during Ramadan. *MCN Am J Matern/Child Nurs* 2011;36:216–21.
 - [69] Manerker K, Harding J, Conlon C, McKinlay C. Maternal gestational diabetes and infant feeding, nutrition and growth: a systematic review and meta-analysis. *Br J Nutr* 2020;123:1201–15.
 - [70] Gilavand A. Studying effect of fasting during Ramadan on mental health of University students in Iran: A review. *J Res Med Dent Sci* 2018;6:205–9.
 - [71] Bandarian F, Namazi N, Atlasi R, Nasli-Esfahani E, Larjani B. Research gaps in Ramadan fasting studies in health and disease. *Diabetes Metabol Syndr: Clin Res Rev* 2021;15:831–5.
 - [72] Rubio-Aliaga I, De Roos B, Duthie SJ, Crosley LK, Mayer C, Horgan G, Colquhoun IJ, Le Gall G, Huber F, Kremer W, et al. Metabolomics of prolonged fasting in humans reveals new catabolic markers. *Metabolomics* 2011;7: 375–87.
 - [73] Mathew S, Krug S, Skurk T, Halama A, Stank A, Artati A, Prehn C, Malek JA, Kastenmüller G, Römisch-Margl W, et al. Metabolomics of Ramadan fasting: an opportunity for the controlled study of physiological responses to food intake. *J Transl Med* 2014;12: 161–161.
 - [74] Harney DJ, Hutchison AT, Hatchwell L, Humphrey SJ, James DE, Hocking S, Heilbronn LK, Larance M. Proteomic analysis of human plasma during intermittent fasting. *J Proteome Res* 2019;18:2228–40.
 - [75] Mindikoglu AL, Abdulsada MM, Jain A, Jalal PK, Devaraj S, Wilhelm ZR, Opekun AR, Jung SY. Intermittent fasting from dawn to sunset for four consecutive weeks induces anticancer serum proteome response and improves metabolic syndrome. *Sci Rep* 2020;10.
 - [76] Bull MJ, Plummer NT. Part 1: the human gut microbiome in health and disease. *Integr Med* 2014;13:17–22.
 - [77] Kho ZY, Lal SK. The human gut microbiome – a potential controller of wellness and disease. *Front Microbiol* 2018;9: 1835–1835.
 - [78] Ali I, Liu K, Long D, Faisal S, Hilal MG, Ali I, Huang X, Long R. Ramadan fasting leads to shifts in human gut microbiota structured by dietary composition. *Front Microbiol* 2021;12.
 - [79] Su J, Wang Y, Zhang X, Ma M, Xie Z, Pan Q, Ma Z, Peppelenbosch MP. Remodeling of the gut microbiome during Ramadan-associated intermittent fasting. *Am J Clin Nutr* 2021;113:1332–42.
 - [80] Ozkul C, Yalinay M, Karaka T. Structural changes in gut microbiome after Ramadan fasting: a pilot study. *Benef Microbes* 2022;227–233.
 - [81] Zeb F, Wu X, Chen L, Fatima S, Ijaz UI H, Chen A, Xu C, Jianglei R, Feng Q, Li M. Time-restricted feeding is associated with changes in human gut microbiota related to nutrient intake. *Nutrition* 2020;78:110797.
 - [82] Elhadd T, Mall R, Bashir M, Palotti J, Fernandez-Luque L, Farooq F, Mohanadi DA, Dabbous Z, Malik RA, Abou-Samra AB. Artificial Intelligence (AI) based machine learning models predict glucose variability and hypoglycaemia risk in patients with type 2 diabetes on a multiple drug regimen who fast during Ramadan (The PROFAST – IT Ramadan study). *Diabetes Res Clin Pract* 2020;169:108388.
 - [83] Hannan MA, Rahman MA, Rahman MS, Sohag AAM, Dash R, Hossain KS, Farjana M, Uddin MJ. Intermittent fasting, a possible priming tool for host defense against SARS-CoV-2 infection: crosstalk among calorie restriction, autophagy and immune response. *Immunol Lett* 2020;226:38–45.
 - [84] Gnoni M, Beas R, Vázquez-Garagatti R. Is there any role of intermittent fasting in the prevention and improving clinical outcomes of COVID-19? intersection between inflammation, mTOR pathway, autophagy and calorie restriction. *VirusDisease* 2021;32:625–34.
 - [85] Ealey KN, Phillips J, Sung H-K. COVID-19 and obesity: fighting two pandemics with intermittent fasting. *Trends Endocrinol Metabol* 2021;32: 706–20.
 - [86] Faris ME, Salem ML, Jahrami HA, Madkour MI, BaHammam AS. Ramadan intermittent fasting and immunity: an important topic in the era of COVID-19. *Ann Thorac Med* 2020;15:125–33.
 - [87] Moghadam MT, Taati B, Paydar Ardakani SM, Suzuki K. Ramadan fasting during the COVID-19 pandemic; observance of health, nutrition and exercise criteria for improving the immune system. *Front Nutr* 2021;7.
 - [88] Bhatti SI, Mindikoglu AL. The impact of dawn to sunset fasting on immune system and its clinical significance in COVID-19 pandemic. *Metabol Open* 2022;13:100162.
 - [89] Akbari HA, Ghram A, Yoosefi M, Arena R, Lavie CJ, Chtourou H, Saad HB, Chamari K. The COVID-19 pandemic and physical activity during intermittent fasting, is it safe? A call for action. *Biol Sport* 2021;38:729–32.
 - [90] Martín-Martín A, Thelwall M, Orduna-Malea E, Delgado López-Cózar E, Google Scholar. Microsoft academic, Scopus, Dimensions, web of science, and OpenCitations' COCI: a multidisciplinary comparison of coverage via citations. *Scientometrics* 2021;126:871–906.
 - [91] Visser M, Van Eck NJ, Waltman L. Large-scale comparison of bibliographic data sources: Scopus, web of science, Dimensions, crossref, and Microsoft academic. *Quan Sci Stud* 2021;2:20–41.
 - [92] Germanis A, Kokkinides P, Stavropoulos-Giokas C. Non-indexed medical journals in the Web: new perspectives in the medical literature. *Int J Med Inf* 1997;47:65–8.
 - [93] Beshyah SA. Authors' selection of target journals and their attitudes to emerging journals: a survey from two developing regions. *Sultan Qaboos University Medical Journal* 2019;19:e51.
 - [94] Trepanowski JF, Bloomer RJ. The impact of religious fasting on human health. *Nutr J* 2010;9:57.
 - [95] El Ati J, Beji C, Danguir J. Increased fat oxidation during Ramadan fasting in healthy women: an adaptive mechanism for body-weight maintenance. *Am J Clin Nutr* 1995;62:302–7.
 - [96] Al-Arouj M, Bouguerra R, Buse J, Hafez S, Hassanein M, Ibrahim MA, Ismail-Beigi F, El-Kebbi I, Khatib O, Kishawi S, et al. Recommendations for management of diabetes during Ramadan. *Diabetes Care* 2005;28:2305–11.
 - [97] Leiper JB, Molla AM, Molla AM. Effects on health of fluid restriction during fasting in Ramadan. *Eur J Clin Nutr* 2003;57(Suppl 2):S30–8.
 - [98] Bogdan A, Bouchareb B, Touitou Y. Ramadan fasting alters endocrine and neuroendocrine circadian patterns. Meal-time as a synchronizer in humans? *Life Sci* 2001;68:1607–15.
 - [99] Adlouni A, Ghalim N, Benslimane A, Lecerf JM, Saile R. Fasting during Ramadan induces a marked increase in high-density lipoprotein cholesterol and decrease in low-density lipoprotein cholesterol. *Ann Nutr Metab* 1997;41:242–9.
 - [100] Bravis V, Hui E, Salih S, Mehara S, Hassanein M, Devendra D. Ramadan education and awareness in diabetes (READ) programme for Muslims with type 2 diabetes who fast during Ramadan. *Diabet Med* 2010;27:327–31.
 - [101] Roky R, Houti I, Moussamih S, Qotbi S, Aadil N. Physiological and

- chronobiological changes during Ramadan intermittent fasting. *Ann Nutr Metab* 2004;48:296–303.
- [103] Hassanein M, Al-Arouj M, Hamdy O, Bebakar WMW, Jabbar A, Al-Madani A, Hanif W, Lessan N, Basit A, Tayeb K, et al. Diabetes and ramadan: practical guidelines. *Diabetes Res Clin Pract* 2017;126:303–16.
- [104] Roky R, Chapotot F, Hakkou F, Bencheikroun MT, Buguet A. Sleep during Ramadan intermittent fasting. *J Sleep Res* 2001;10:319–27.
- [105] Sweileh N, Schnitzler A, Hunter GR, Davis B. Body composition and energy metabolism in resting and exercising Muslims during Ramadan fast. *J Sports Med Phys Fit* 1992;32:156–63.
- [106] Iraki L, Bogdan A, Hakkou F, Amrani N, Abkari A, Touitou Y. Ramadan diet restrictions modify the circadian time structure in humans. A study on plasma gastrin, insulin, glucose, and calcium and on gastric pH. *J Clin Endocrinol Metab* 1997;82:1261–73.
- [107] Nag OS. Muslim population by country. 2019.
- [108] Fairs ME, et al. Intermittent fasting during Ramadan is associated with improved plasma sphingolipids and inflammatory markers in subjects with overweight/obese: A lipidomics approach prospective cohort study. 2022. In preparation.