



Changes in Placental Size during Ramadan

S.H. Alwasel^{a,*}, Z. Abotalib^b, J.S. Aljarallah^c, C. Osmond^d, S.M. Alkharaz^e, I.M. Alhazza^a, G. Badr^a, D.J.P. Barker^{a,c,d,f}

^a Fetal Programming of Diseases Research Chair, Zoology Department, College of Science, King Saud University, Saudi Arabia

^b Obstetrics and Gynecology Department, King Saud University, Saudi Arabia

^c Family and Community Medicine Department, King Saud University, Saudi Arabia

^d MRC Epidemiology Resource Centre, Medical School, University of Southampton, UK

^e King Saud Hospital, Unizah, Saudi Arabia

^f Heart Research Center, Oregon Health & Science University, USA

ARTICLE INFO

Article history:

Accepted 21 April 2010

Keywords:

Placental weight

Placental ratio

Ramadan

Programming

ABSTRACT

Background: Placental growth responds to maternal influences. Ramadan is an annual period of day-time fasting during which people in Saudi Arabia, including pregnant women, change their diets and physical activity. Little is known about the effects of this altered lifestyle on placental development.

Methods: We studied the birth records of 7083 babies born over a four-year period to Saudi nationals in Unizah, a small city 350 km to the north of Riyadh, the capital city of Saudi Arabia. The records included birth weight, placental weight and gestational age.

Results: Mean birth weight was similar to European values but the mean placental weight and ratio of placental weight to birth weight were lower. Among babies who were in the second or third trimester of gestation during Ramadan the mean placental weight and ratio were below those of babies who were not in utero during Ramadan. Among boys the mean placental ratios were 14.4 percent (second trimester) and 14.5 percent (third trimester) compared with 14.9 percent ($p = <0.001$ and 0.002). The corresponding figures for girls were 14.8 and 14.6 percent compared with 15.1 percent ($p = 0.02$ and <0.001). **Conclusions:** In Saudi Arabia placentas respond to mothers' limited ability to deliver nutrients to them. Placental growth slows but efficiency is increased so that fetal growth is sustained, albeit with a reduced reserve capacity. The lifestyle changes associated with Ramadan further slow placental growth. Ramadan may influence placental growth through dietary changes other than day-time fasting. Changes in placental growth during Ramadan could be associated with altered fetal programming, and may therefore have long-term implications for the health of the next generation.

© 2010 Elsevier Ltd. All rights reserved.

1. Introduction

In humans, placental growth responds to maternal influences. Maternal anemia and high maternal body mass index are associated with a high placental ratio at birth (placental weight/birth weight) [1,2]. Maternal smoking reduces both placental weight and birth weight. The diets of mothers during pregnancy, and their physical activity, are also known to be associated with altered placental size [2,3]. In sheep farming manipulation of placental size

by moving pregnant ewes to different pastures is standard practice, because it leads to the production of larger lambs [4,5]. There is evidence for similar effects in humans [6].

In Islam, Ramadan is an annual period of day-time fasting. It lasts for one month and occurs at different seasons in different years because the Arabic calendar depends on the moon. Therefore the hours of daily fasting will depend on the season. During Ramadan people in Saudi Arabia change their lifestyle. They take no food or water from dawn to sunset, when they break their fast by eating sweet and fried meals. The next meal is "Sahoor" which is usually eaten before dawn and comprises fat-rich foods. People reduce their activities during the day, but are more active at night. Studies have shown that body weight and body mass index (weight/height²) are reduced during Ramadan in both men and women [7,8]. This may reflect the changes in diet and pattern of eating rather than the

* Corresponding author. Zoology Department, College of Science, King Saud University, PO Box 2455, Riyadh 11451, Saudi Arabia. Tel.: +966 550880848; fax: +966 14678514.

E-mail address: sawasel10@hotmail.com (S.H. Alwasel).

effects of fasting. Although pregnant women are allowed to defer fasting until after the pregnancy in Saudi Arabia they usually prefer to share the spiritual and social experiences of Ramadan with their families. Little is known about the effects of the altered lifestyle during Ramadan on placental development.

Fetal and placental size at birth are associated with the later risk of chronic diseases. People who were born at term but whose birth weights were towards the lower end of the normal range are at increased risk of a range of disorders including coronary heart disease, stroke, hypertension and type 2 diabetes [9,10]. Placental weight correlates with birth weight [11] and is also associated with later cardiovascular disease. Associations between both low placental weight and with high placental weight in relation to birth weight have been reported [2,12–14]. These associations between birth size and later chronic disease are thought to reflect “fetal programming”, the process whereby nutrition and other influences during gestation permanently change the structure and function of the body in ways that affect long-term health [9]. If the altered lifestyle during Ramadan changes fetal or placental growth it could change fetal programming, and could therefore have long-term implications for the health of the next generation. There are sex differences in fetal and placental growth [15,16]. Boys grow more rapidly than girls [17]. At any placental weight at birth boys tend to be longer than girls [18]. This suggests that boy's placentas are more efficient, but they may have less reserve capacity. We present an analysis of body and placental size at birth in 7000 newborn babies who were in utero, at different stages of gestation, during Ramadan. Boys and girls are analysed separately.

2. Methods

We studied the birth records of 7083 babies born in King Saud Hospital, Unizah, Saudi Arabia. Unizah is a small city 350 km to the north of the capital city, Riyadh. It is an agricultural and administrative centre with a population 140,000. The King Saud Hospital is one of two maternity centres in the city. We restricted our data collection to singletons born to Saudi nationals. These were the only exclusion criteria. The data were abstracted from the maternity log books. During the four-year period of our study Ramadan occurred in three different calendar months, from October to December. We calculated whether each baby was in utero during Ramadan and, if so, during which trimester. The hospital records included date of birth, maternal age, birth weight, placental weight and gestational age, estimated from the date of the last menstrual period. Pregnant women are first seen around 40 days after their expected menstrual period did not occur. Women in Saudi Arabia are careful in knowing their menstrual cycles as menstrual bleeding affects religious practices. The majority of the mothers were born in Unizah.

2.1. Statistical methods

We studied births that occurred after 37 completed weeks of gestation. We defined four groups of exposure to Ramadan using the month of birth in the Arabic calendar. For example, births in

months one to three (Muharram, Safar and Rabi' al-awwal) were treated as exposed to Ramadan in the second trimester. We analysed birth weight, placental weight and the ratio of placental to birth weight using multiple linear regression, adjusting for gender, time through the study and gestational age by including them as predictors. We included in the regression dummy variables for the trimester of exposure to Ramadan to assess trimester-specific effects, treating the group not exposed to Ramadan in pregnancy as the baseline group for comparison.

3. Results

There were 7083 birth records. Placental weight was correlated with birth weight ($r = 0.34$) and with gestational age ($r = 0.20$). The mean age of the mothers was 29.5 years. Table 1 shows the mean birth weight, placental weight and gestational age for boys and girls. Boys had higher mean birth weights and placental weights than girls but the ratio of placental to birth weight was lower. We found that while babies' mean birth weights remained constant throughout the four years of the study, their mean placental weights increased by 29 g per year, so that the ratio of placental weight to birth weight rose. We therefore adjusted for year of birth in our analyses.

In Table 2 the boys and girls are divided according to whether or not they were in utero during Ramadan and if so, whether it was during the first, second or third trimester of gestation. Among babies who were in the first trimester of gestation during Ramadan there was no difference in birth weight or placental size compared with babies who were not in utero during Ramadan. Among babies who were in the second or third trimester during Ramadan the mean birth weight was the same as that of babies who were not in utero during Ramadan, but mean placental weight and the ratio of placental weight to birth weight were lower. These differences were little changed by adjusting for the length of gestation.

Figure 1 shows that, while mean placental weights were higher in boys than girls, in both boys and girls those in the second or third trimester of gestation during Ramadan had reduced mean values. Similarly Figure 2 shows that, while the ratios of placental weight to birth weight were lower in boys than girls irrespective of whether or not they were in utero during Ramadan, in both boys and girls those in the second or third trimester of gestation during Ramadan had reduced mean ratios.

4. Discussion

We have examined the body and placental size of 7000 babies born over a four-year period in a small city in Saudi Arabia. When compared to babies who were not in utero during Ramadan those who were in the second or third trimester of gestation during Ramadan had reduced mean placental weight and a reduced ratio of placental weight to birth weight. The occurrence of Ramadan was not associated with changes in birth weight.

Placental weight was correlated with both birth weight and gestational age, as would be expected. In contrast to mean birth

Table 1
Mean birth size and gestational age in boys and girls.

Measurements	Boys		Girls		p-Value for difference
	Mean (SD)	n	Mean (SD)	n	
Birth weight (kg)	3.29 (0.50)	3542	3.19 (0.49)	3541	<0.001
Placental weight (g)	473 (87)	3498	467 (86)	3509	0.002
Placental/birth weight (%)	14.6 (3.1)	3498	14.9 (3.0)	3509	<0.001
Gestational age (weeks)	39.8 (1.1)	3542	39.8 (1.0)	3541	0.7

Table 2

Mean birth size (SD) according to timing of Ramadan during pregnancy.

Measurement	Timing of Ramadan during pregnancy					
	Outside of pregnancy	First trimester	p^*	Second trimester	p^*	Third trimester
Boys number	904	771		976		891
Birth weight (kg)	3.28 (0.51)	3.31 (0.50)	0.2	3.29 (0.47)	0.7	3.30 (0.50)
Placental weight (g)	479 (81)	481 (82)	0.8	466 (78)	<0.001	468 (80)
Placental/birth weight (%)	14.9 (2.9)	14.8 (3.0)	0.4	14.4 (2.8)	<0.001	14.5 (3.0)
Girls number	918	827		958		838
Birth weight (kg)	3.17 (0.48)	3.20 (0.49)	0.3	3.18 (0.48)	0.7	3.20 (0.48)
Placental weight (g)	471 (75)	472 (83)	0.9	463 (79)	0.02	460 (82)
Placental/birth weight (%)	15.1 (2.8)	15.0 (3.0)	0.6	14.8 (2.8)	0.02	14.6 (2.8)

 $p^* = p$ Value for comparison with babies not in utero during Ramadan.

weight, which approached the mean for western populations, the mean placental weights and placental to birth weight ratios (Table 1) were well below those in a recent compilation of European reference values [19], and were also below those recorded in Europe before the second world war [6]. One possible explanation is that the techniques used for trimming and weighing the placenta were different in our study and produced a systematic error. We have inquired about the techniques used in the King Saud Hospital and they are similar to those used in Europe and the United States. Measurements of placental weight during routine clinical practice are subject to random errors, which may be greater than those associated with birth weight. In Saudi Arabia, however, the placenta is handled carefully: it remains valued after birth and is taken to a mortuary for burial. We therefore conclude that the low placental weights recorded in our study reflect small placental size in relation to western populations. Measurement error is unlikely to explain the differences in placental size among babies who were or were not in utero during Ramadan.

In experimental studies in which uterine blood flow is reduced [20], and in clinical studies [21], a relative decrease in placental size is evident before changes in fetal growth. The placenta seems to act as a nutrient sensor regulating the transfer of nutrients to the fetus according to the mother's ability to deliver them [3]. The growth of every human fetus may be constrained by the limited capacity of the mother to deliver nutrients to it, and this capacity may originate in the mother's own fetal life [22]. The placenta's ability to transfer nutrients is reflected in its size: small babies generally have small placentas [11]. An interpretation of our findings is that in Saudi Arabia placentas respond to mothers' limited ability to deliver nutrients to them. Placental growth slows but efficiency is increased

so that fetal growth is sustained, albeit with a reduced reserve capacity. We do not have any evidence that this reduced reserve affects pregnancy outcome. The lifestyle changes associated with Ramadan further slow placental growth with a further increase in efficiency. The increase in placental size and in the ratio of placental weight to birth weight over the four years of our study could indicate that maternal nutrient delivery in Saudi Arabia has improved.

In our study boys had higher mean birth and placental weights than girls, but a lower ratio of placental weight to birth weight. This is consistent with other findings [17,19]. It has led to the conclusion that boy's placentas are more efficient than girl's placentas but may have a lesser reserve capacity [17]. Ramadan was associated with similar changes in placental size in boys and girls (Figures 1 and 2). In the war-time famine in Holland fewer boys than girls were born alive [23], suggesting that boys are more dependent on the mother's food intake, as opposed to her food stores and metabolism [17]. The lack of a selective effect of Ramadan on boys may be evidence that it influences placental growth through dietary changes other than day-time fasting.

The delivery of nutrients from the mother can alter fetoplacental growth and programme the fetus from the earliest stages of development, even before implantation [24]. The placentas of babies who were in the first trimester of gestation during Ramadan, however, were similar to those who were not in utero during Ramadan. This implies that dietary changes during Ramadan influence placental growth once the maternal blood supply is established and maternal blood enters the placenta.




Fig. 1. Mean placental weight according to time in pregnancy when Ramadan occurred. $p = p$ value for comparison with babies not in utero during Ramadan.




Fig. 2. Mean ratio of placental weight to birth weight according to time in pregnancy when Ramadan occurred. $p = p$ value for comparison with babies not in utero during Ramadan.

4.1. Limitations of the study

Our study was based on measurements of placental and birth weights made during routine clinical practice. We have discussed the possible measurement errors. Our study sample came from one city in Saudi Arabia and may not represent the country as a whole. We collected data over a period of four years during which Ramadan occurred in only three different months. Therefore we cannot distinguish the effects of Ramadan from seasonal effects on food intake or the pattern of eating. Our data came from log books kept in the labour ward. We did not have access to the patients' case records and were therefore unable to explore whether the changes in placental weight at birth according to the timing of Ramadan during gestation were associated with changes in maternal weight, weight gain or haemoglobin. Neither were we able to examine the effects of parity on placental weight. Our measurements of the length of gestation were based on the date of the mother's last menstrual period and would therefore be subject to error.

5. Conclusion

We have found that babies who were in the second or third trimester of gestation during the Ramadan fasts had reduced placental weight at birth. Birth weight was not affected, which suggests that the placentas were able to maintain levels of activity despite their reduced size. The placental weights recorded in our study were small in relation to those in western populations whereas the birth weights were similar. An interpretation of our findings is that in Saudi Arabia placentas respond to mothers' limited ability to deliver nutrients to them. Their growth is slowed but their efficiency is increased so that fetal growth is sustained, albeit with a reduced reserve capacity. The lifestyle changes associated with Ramadan further slow placental growth, with a further increase in efficiency. Changes in placental growth during Ramadan could be associated with altered fetal programming, and may therefore have long-term implications for the health of the next generation.

References

- [1] Godfrey KM, Redman CWG, Barker DJP, Osmond C. The effect of maternal anaemia and iron deficiency on the ratio of fetal weight to placental weight. *Br J Obstet Gynaecol* 1991;98:886–91.
- [2] Godfrey KM. The role of the placenta in fetal programming. *Placenta* 2002;23 (Suppl A):S20–7.
- [3] Jansson T, Powell TL. Role of the placenta in fetal programming: underlying mechanisms and potential interventional approaches. *Clin Sci* 2007;113:1–13.
- [4] McCrabb GJ, Egan AR, Hosking BJ. Maternal undernutrition during mid-pregnancy in sheep. Placental size and its relationship to calcium transfer during late pregnancy. *Br J Nutr* 1991;65:157–68.
- [5] McCrabb GJ, Egan AR, Hosking BJ. Maternal undernutrition during mid-pregnancy in sheep: variable effects on placental growth. *J Agric Sci* 1992;118:127–32.
- [6] Barker DJP, Thornburg K, Osmond C, Kajantie E, Eriksson JG. The surface area of the placenta and hypertension in the offspring in later life. *Int J Dev Biol* 2010;54:525–30.
- [7] Ziaee V, Razaei M, Ahmadinejad Z, Shaikh H, Yousefi R, Yarmohammadi L, et al. The changes of metabolic profile and weight during Ramadan fasting. *Singapore Med J* 2006;47(5):409–14.
- [8] Lamri-Senhadj SY, El Kebir B, Belleville J, Bouchenak M. Assessment of dietary consumption and time course changes in serum lipids and lipoproteins before, during and after Ramadan in young Algerian adults. *Singapore Med J* 2009;50 (3):288–94.
- [9] Barker DJP. Fetal origins of coronary heart disease. *Br Med J* 1995;311:171–4.
- [10] Barker DJP, Osmond C, Kajantie E, Eriksson JG. Growth and chronic disease: findings in the Helsinki Birth Cohort. *Ann Hum Biol* 2009;36:445–58.
- [11] Sibley CP. In: Case RM, Waterhouse JM, editors. *Human physiology: age, stress, and the environment*. Oxford: Oxford University Press; 1994. p. 3–27.
- [12] Barker DJP, Bull AR, Osmond C, Simmonds S. Fetal and placental size and risk of hypertension in adult life. *Br Med J* 1990;301:259–62.
- [13] Campbell DM, Hall MH, Barker DJP, Cross J, Shiell A W, Godfrey KM. Diet in pregnancy and the offspring's blood pressure 40 years later. *Brit J Obstet Gynaecol* 1996;103:273–80.
- [14] Eriksson J, Forsen T, Tuomilehto J, Osmond C, Barker D. Fetal and childhood growth and hypertension in adult life. *Hypertension* 2000;36:790–4.
- [15] Pedersen JF. Ultrasound evidence of sexual difference in fetal size in first trimester. *Br Med J* 1980;281:1253.
- [16] Tanner JM. *Fetus into man*. 2nd ed. Ware: Castlemead; 1989.
- [17] Eriksson JG, Kajantie E, Osmond C, Thornburg K, Barker DJP. Boys live dangerously in the womb. *Am J Human Biol*; 2010. doi:10.1002/ajhb.20995.
- [18] Forsén T, Eriksson JG, Tuomilehto J, Osmond C, Barker DJP. Growth in utero and during childhood among women who develop coronary heart disease: longitudinal study. *Br Med J* 1999;319:1403–7.
- [19] Burkhard T, Schaffer L, Schneider S, Zimmermann R, Kurmanavicius J. Reference values for the weight of freshly delivered term placentas and for placental weight–birth weight ratios. *Eur J Obstet Gynaecol* 2006;128:248–52.
- [20] Jansson T, Thordstein M, Kjellmer I. Placental blood flow and fetal weight following uterine artery ligation. *Biol Neonate* 1986;49:172–80.
- [21] Hafner E, Metzenbauer M, Hofinger D, et al. Placental growth from the first to the second trimester of pregnancy in SGA-foetuses and pre-eclamptic pregnancies compared to normal foetuses. *Placenta* 2003;24:336–42.
- [22] Ounsted M, Scott A, Ounsted C. Transmission through the female line of a mechanism constraining human fetal growth. *Ann Hum Biol* 1986;13:143–51.
- [23] Ravelli AC, van der Meulen JHP, Osmond C, Barker DJP, Bleker OP. Obesity at the age of 50 years in men and women exposed to famine prenatally. *Am J Clin Nutr* 1999;70:811–6.
- [24] Kwong WY, Wild AE, Roberts P, Willis AC, Fleming TP. Maternal undernutrition during the preimplantation period of rat development causes blastocyst abnormalities and programming of postnatal hypertension. *Development* 2000;127:4195–202.